

EVALUATION OF n + 54Fe CROSS SECTIONS FOR THE ENERGY  
RANGE 1.0E-11 to 150 MeV

M. B. Chadwick, P. G. Young, and A. J. Koning  
6 September 1996

This evaluation provides a complete representation of the nuclear data needed for transport, damage, heating, radioactivity, and shielding applications over the incident neutron energy range from 1.0E-11 to 150 MeV. The discussion here is divided into the region below and above 20 Mev.

INCIDENT NEUTRON ENERGIES < 20 MeV

Below 20 MeV the evaluation is based completely on the ENDF/B-VI.1 (Release 1) evaluation by He91 (see also Fu86). The following modifications were made to the ENDF/B-VI.1 evaluation:

1. The covariance files (MF=33) were removed from the file.
2. The derived MF=3 files for MT=203,205,207 were removed.

INCIDENT NEUTRON ENERGIES > 20 MeV

The evaluation above 20 MeV utilizes MF=6, MT=5 to represent all reaction data. Production cross sections and emission spectra are given for neutrons, protons, deuterons, tritons, alpha particles, gamma rays, and all residual nuclides produced ( $A>5$ ) in the reaction chains. To summarize, the ENDF sections with non-zero data above  $En = 20$  MeV are:

MF=3 MT= 1 Total Cross Section  
MT= 2 Elastic Scattering Cross Section  
MT= 3 Nonelastic Cross Section  
MT= 5 Sum of Binary (n,n') and (n,x) Reactions

MF=4 MT= 2 Elastic Angular Distributions

MF=6 MT= 5 Production Cross Sections and Energy-Angle Distributions for Emission Neutrons, Protons, Deuterons, and Alphas; and Angle-Integrated Spectra for Gamma Rays and Residual Nuclei That Are Stable Against Particle Emission

The evaluation is based on nuclear model calculations that have been benchmarked to experimental data, especially for n + Fe56 and p + Fe56 reactions (Ch96a). We use the GNASH code system (Yo92), which utilizes Hauser-Feshbach statistical, preequilibrium and direct-reaction theories. Spherical optical model calculations are used to obtain particle transmission coefficients for the Hauser-Feshbach calculations, as well as for the elastic neutron angular distributions.

Cross sections and spectra for producing individual residual nuclei are included for reactions that exceed a cross section of approximately 1 nb at any energy. The energy-angle-correlations for all outgoing particles are based on Kalbach systematics (Ka88).

A model was developed to calculate the energy distributions of

all recoil nuclei in the GNASH calculations (Ch96b). The recoil energy distributions are represented in the laboratory system in MT=5, MF=6, and are given as isotropic in the lab system. Note that all other data in MT=5, MF=6 are given in the center-of-mass system. This method of representation requires a modification of the original ENDF-6 format.

Preequilibrium corrections were performed in the course of the GNASH calculations using the exciton model of Kalbach (Ka77, Ka85), validated by comparison with calculations using Feshbach, Kerman, Koonin (FKK) theory [Ch93]. Discrete level data from nuclear data sheets were matched to continuum level densities using the formulation of Ignatyuk (Ig75) and pairing and shell parameters from the Cook (Co67) analysis. Neutron and charged-particle transmission coefficients were obtained from the optical potentials, as discussed below. Gamma-ray transmission coefficients were calculated using the Kopecky-Uhl model (Ko90).

The neutron optical model potential of Arthur et al. (Ar80) was used to calculate transmission coefficients and cross sections with the SCAT2 code (Be92) up to a neutron energy of 26 MeV. Between 26 and 52 MeV, the imaginary volume component of Arthur's potential was modified to better account for nonelastic cross section measurements, and above 52 MeV the Semmering potential of Madland (Ma88) was used. For protons, the Beccetti-Greenlees potential (Be69) was utilized below 28 MeV, and the Madland potential (Ma88) was used at higher energies. The global spherical potential of Perey (Pe63) was utilized for deuterons, and the potential of Beccetti-Greenlees (Be) was adopted for tritons. Finally, the alpha potential of Lemos (Le72), as adapted by Arthur et al. (Ar80), was used for alpha particles.

Direct reaction cross sections to discrete states were calculated with the ECIS95 code [Re95] using deformation parameters compiled in Nuclear Data Sheets.

We used the same values for the total, elastic, and nonelastic cross sections (MF=3, MT=1,2,3) above 20 MeV as were used in the n + 56Fe evaluation.

\*\*\*\*\*

#### REFERENCES

- [Ar80]. E.D. Arthur and P.G. Young, 'Evaluation of Neutron Cross Sections to 40 MeV for 54,56Fe," Proc. Sym. on Neutron Cross Sections from 10 to 50 MeV, 12-14 May 1980, Brookhaven National Laboratory [Eds. M. R. Bhat and S. Pearlstein, BNL-NCS-51245, 1980] p. 731.
- [Be69]. F.D. Becchetti, Jr., and G.W. Greenlees, Phys. Rev. 182, 1190 (1969).
- [Be71]. F.D. Becchetti, Jr., and G.W. Greenlees in "Polarization Phenomena in Nuclear Reactions," (Ed: H.H. Barschall and W. Haeberli, The University of Wisconsin Press, 1971) p.682.
- [Be92]. O. Bersillon, "SCAT2 - A Spherical Optical Model Code," in Proc. ICTP Workshop on Computation and Analysis of Nuclear Data Relevant to Nuclear Energy and Safety, 10 February-13 March, 1992, Trieste, Italy, to be published in World Scientific Press, and Progress Report of the Nuclear Physics Division,

Bruyeres-le-Chatel 1977, CEA-N-2037, p.111 (1978).

[Ch93]. M. B. Chadwick and P. G. Young, "Feshbach-Kerman-Koonin Analysis of  $^{93}\text{Nb}$  Reactions:  $P \rightarrow Q$  Transitions and Reduced Importance of Multistep Compound Emission," Phys. Rev. C 47, 2255 (1993).

[Ch96a]. M. B. Chadwick and P. G. Young, "GNASH Calculations of  $n, p + 54, 56, 57, 58\text{Fe}$  and Benchmarking of Results" in APT PROGRESS REPORT: 1 August - 1 September 1996, internal Los Alamos National Laboratory memo T-2-96/MS-52, 6 Aug. 1996 from R.E. MacFarlane to L. Waters.

[Ch96b]. M. B. Chadwick, P. G. Young, R. E. MacFarlane, and A. J. Koning, "High-Energy Nuclear Data Libraries for Accelerator-Driven Technologies: Calculational Method for Heavy Recoils," Proc. of 2nd Int. Conf. on Accelerator Driven Transmutation Technology and Applications, Kalmar, Sweden, 3-7 June 1996.

[Co67]. J. L. Cook, H. Ferguson, and A. R. Musgrove, "Nuclear Level Densities in Intermediate and Heavy Nuclei," Aust.J.Phys. 20, 477 (1967).

[Fu86]. C. Y. Fu and D. M. Hetrick, ORNL/TM-9964, ENDF-341 (1986).

[He91]. D. M. Hetrick, C. Y. Fu, and N. M. Larson, ENDF/B-VI.1 Evaluation of  $n + 54\text{Fe}$ , personal comm., (1991).

[Ig75]. A. V. Ignatyuk, G. N. Smirenkin, and A. S. Tishin, "Phenomenological Description of the Energy Dependence of the Level Density Parameter," Sov. J. Nucl. Phys. 21, 255 (1975).

[Ka77]. C. Kalbach, "The Griffin Model, Complex Particles and Direct Nuclear Reactions," Z.Phys.A 283, 401 (1977).

[Ka85]. C. Kalbach, "PRECO-D2: Program for Calculating Preequilibrium and Direct Reaction Double Differential Cross Sections," Los Alamos National Laboratory report LA-10248-MS (1985).

[Ka88]. C. Kalbach, "Systematics of Continuum Angular Distributions: Extensions to Higher Energies," Phys.Rev.C 37, 2350 (1988); see also C. Kalbach and F. M. Mann, "Phenomenology of Continuum Angular Distributions. I. Systematics and Parameterization," Phys.Rev.C 23, 112 (1981).

[Ko90]. J. Kopecky and M. Uhl, "Test of Gamma-Ray Strength Functions in Nuclear Reaction Model Calculations," Phys.Rev.C 42, 1941 (1990).

[Le72]. O. F. Lemos, "Diffusion Elastique de Particules Alpha de 21 à 29.6 MeV sur des Noyaux de la Région Ti-Zn," Orsay report, Series A, No. 136 (1976).

[Ma88]. D.G. Madland, "Recent Results in the Development of a Global Medium-Energy Nucleon-Nucleus Optical-Model Potential," Proc. OECD/NEANDC Specialist's Mtg. on Preequilibrium Nuclear Reactions, Semmering, Austria, 10-12 Feb. 1988, NEANDC-245 'U' (1988).

[Pe63]. C. M. Perey and F. G. Perey, Phys. Rev. 132, 755 (1963).

[Re95]. J. Raynal, "Notes on ECIS94," CEA informal report, Saclay (1995).

[Yo92]. P. G. Young, E. D. Arthur, and M. B. Chadwick, "Comprehensive Nuclear Model Calculations: Introduction to the Theory and Use of the GNASH Code," LA-12343-MS (1992).

26054 = TARGET 1000Z+A (if A=0 then elemental)

1 = PROJECTILE 1000Z+A

Nonelastic, elastic, and Production cross sections for A&lt;5 projectiles in barns:

Energy	nonelas	elastic	neutron	proton	deuteron	triton	helium3	alpha	gamma
2.000E+01	1.296E+00	9.424E-01	1.188E+00	9.654E-01	3.292E-02	7.686E-04	0.000E+00	7.651E-02	2.380E+00
2.200E+01	1.240E+00	9.979E-01	1.149E+00	9.822E-01	3.755E-02	1.319E-03	0.000E+00	8.006E-02	2.434E+00
2.400E+01	1.177E+00	1.053E+00	1.102E+00	1.018E+00	4.024E-02	1.927E-03	0.000E+00	8.035E-02	2.351E+00
2.600E+01	1.117E+00	1.135E+00	1.086E+00	1.027E+00	4.459E-02	2.507E-03	0.000E+00	7.915E-02	2.209E+00
2.800E+01	1.084E+00	1.206E+00	1.116E+00	1.029E+00	4.928E-02	3.137E-03	0.000E+00	7.756E-02	2.119E+00
3.000E+01	1.063E+00	1.267E+00	1.151E+00	1.043E+00	5.366E-02	3.740E-03	0.000E+00	7.630E-02	2.051E+00
3.500E+01	1.012E+00	1.422E+00	1.206E+00	1.070E+00	6.087E-02	4.987E-03	0.000E+00	7.461E-02	1.973E+00
4.000E+01	9.637E-01	1.536E+00	1.234E+00	1.086E+00	6.562E-02	5.827E-03	0.000E+00	7.441E-02	1.880E+00
4.500E+01	9.265E-01	1.593E+00	1.283E+00	1.125E+00	6.886E-02	6.403E-03	0.000E+00	7.656E-02	1.787E+00
5.000E+01	9.004E-01	1.600E+00	1.343E+00	1.180E+00	6.977E-02	6.852E-03	0.000E+00	8.161E-02	1.756E+00
5.500E+01	8.778E-01	1.587E+00	1.407E+00	1.244E+00	7.093E-02	6.916E-03	0.000E+00	8.923E-02	1.725E+00
6.000E+01	8.578E-01	1.547E+00	1.458E+00	1.291E+00	7.235E-02	7.178E-03	0.000E+00	9.573E-02	1.699E+00
6.500E+01	8.399E-01	1.488E+00	1.522E+00	1.347E+00	7.394E-02	7.645E-03	0.000E+00	1.071E-01	1.659E+00
7.000E+01	8.236E-01	1.428E+00	1.556E+00	1.396E+00	7.461E-02	8.117E-03	0.000E+00	1.164E-01	1.556E+00
7.500E+01	8.087E-01	1.352E+00	1.609E+00	1.436E+00	7.538E-02	8.647E-03	0.000E+00	1.258E-01	1.535E+00
8.000E+01	7.949E-01	1.279E+00	1.667E+00	1.477E+00	7.735E-02	9.377E-03	0.000E+00	1.370E-01	1.512E+00
8.500E+01	7.823E-01	1.218E+00	1.737E+00	1.550E+00	8.124E-02	1.033E-02	0.000E+00	1.492E-01	1.466E+00
9.000E+01	7.705E-01	1.159E+00	1.780E+00	1.584E+00	8.331E-02	1.117E-02	0.000E+00	1.587E-01	1.441E+00
9.500E+01	7.596E-01	1.090E+00	1.823E+00	1.611E+00	8.550E-02	1.210E-02	0.000E+00	1.680E-01	1.429E+00
1.000E+02	7.494E-01	1.037E+00	1.865E+00	1.643E+00	8.800E-02	1.329E-02	0.000E+00	1.788E-01	1.405E+00
1.100E+02	7.307E-01	9.113E-01	1.930E+00	1.683E+00	9.232E-02	1.555E-02	0.000E+00	1.959E-01	1.359E+00
1.200E+02	7.141E-01	8.009E-01	2.004E+00	1.723E+00	9.541E-02	1.839E-02	0.000E+00	2.124E-01	1.306E+00
1.300E+02	6.992E-01	7.028E-01	2.059E+00	1.764E+00	9.944E-02	2.125E-02	0.000E+00	2.263E-01	1.252E+00
1.400E+02	6.857E-01	6.243E-01	2.114E+00	1.798E+00	1.043E-01	2.452E-02	0.000E+00	2.396E-01	1.198E+00
1.500E+02	6.734E-01	5.896E-01	2.150E+00	1.823E+00	1.085E-01	2.771E-02	0.000E+00	2.498E-01	1.168E+00


26054 = TARGET 1000Z+A (if A=0 then elemental)

1 = PROJECTILE 1000Z+A







Kerma coefficients in units of f.Gy.m^2:

Energy	proton	deuteron	triton	helium3	alpha	non-rec	elas-rec	TOTAL
2.000E+01	9.635E-01	5.320E-02	6.899E-04	0.000E+00	1.345E-01	1.137E-01	2.039E-02	1.286E+00
2.200E+01	1.002E+00	6.796E-02	1.406E-03	0.000E+00	1.454E-01	1.177E-01	3.311E-02	1.367E+00
2.400E+01	1.050E+00	8.099E-02	2.370E-03	0.000E+00	1.514E-01	1.195E-01	3.564E-02	1.439E+00
2.600E+01	1.081E+00	9.754E-02	3.467E-03	0.000E+00	1.543E-01	1.200E-01	3.862E-02	1.495E+00
2.800E+01	1.124E+00	1.174E-01	4.846E-03	0.000E+00	1.564E-01	1.220E-01	3.929E-02	1.564E+00
3.000E+01	1.181E+00	1.387E-01	6.362E-03	0.000E+00	1.585E-01	1.245E-01	3.889E-02	1.648E+00
3.500E+01	1.341E+00	1.887E-01	1.033E-02	0.000E+00	1.637E-01	1.320E-01	3.667E-02	1.872E+00
4.000E+01	1.504E+00	2.401E-01	1.422E-02	0.000E+00	1.696E-01	1.375E-01	3.370E-02	2.099E+00
4.500E+01	1.678E+00	2.889E-01	1.788E-02	0.000E+00	1.786E-01	1.435E-01	3.064E-02	2.338E+00
5.000E+01	1.878E+00	3.226E-01	2.138E-02	0.000E+00	1.923E-01	1.494E-01	2.764E-02	2.591E+00
5.500E+01	2.101E+00	3.589E-01	2.322E-02	0.000E+00	2.090E-01	1.565E-01	2.289E-02	2.872E+00
6.000E+01	2.304E+00	3.958E-01	2.581E-02	0.000E+00	2.250E-01	1.617E-01	2.093E-02	3.133E+00
6.500E+01	2.492E+00	4.303E-01	2.829E-02	0.000E+00	2.493E-01	1.685E-01	1.915E-02	3.388E+00
7.000E+01	2.685E+00	4.570E-01	3.070E-02	0.000E+00	2.702E-01	1.745E-01	1.767E-02	3.635E+00
7.500E+01	2.864E+00	4.817E-01	3.301E-02	0.000E+00	2.914E-01	1.796E-01	1.621E-02	3.866E+00
8.000E+01	3.038E+00	5.148E-01	3.532E-02	0.000E+00	3.160E-01	1.852E-01	1.496E-02	4.104E+00
8.500E+01	3.296E+00	5.646E-01	3.858E-02	0.000E+00	3.448E-01	1.919E-01	1.396E-02	4.450E+00
9.000E+01	3.475E+00	6.012E-01	4.097E-02	0.000E+00	3.670E-01	1.961E-01	1.309E-02	4.693E+00
9.500E+01	3.655E+00	6.386E-01	4.340E-02	0.000E+00	3.890E-01	1.994E-01	1.215E-02	4.938E+00
1.000E+02	3.827E+00	6.731E-01	4.592E-02	0.000E+00	4.139E-01	2.036E-01	1.144E-02	5.175E+00
1.100E+02	4.169E+00	7.452E-01	5.069E-02	0.000E+00	4.556E-01	2.094E-01	9.904E-03	5.639E+00
1.200E+02	4.512E+00	7.785E-01	5.605E-02	0.000E+00	4.974E-01	2.134E-01	8.615E-03	6.066E+00
1.300E+02	4.854E+00	8.331E-01	6.117E-02	0.000E+00	5.338E-01	2.241E-01	7.503E-03	6.514E+00
1.400E+02	5.176E+00	8.947E-01	6.673E-02	0.000E+00	5.689E-01	2.342E-01	6.624E-03	6.947E+00
1.500E+02	5.504E+00	9.511E-01	7.213E-02	0.000E+00	5.975E-01	2.411E-01	6.220E-03	7.372E+00






$n + {}^{54}\text{Fe}$  nonelastic and production cross sections




# $n + {}^{54}\text{Fe}$ angle-integrated emission spectra



# $n + {}^{54}\text{Fe}$ Kalbach preequilibrium ratios




26-FE-54 APT LA150 NJOY 97.18 MCNPX  
angular distribution for elastic



26-FE-54 APT LA150 NJOY 97.18 MCNPX

Heating



26-FE-54 APT LA150 NJOY 97.18 MCNPX

Damage

